

**EFFECTS OF ZINC SUPPLEMENTATION ON REPRODUCTIVE PERFORMANCE IN CATTLE: A REVIEW****Pengaruh Suplementasi Zink Terhadap Performa Reproduksi Pada Sapi:
A Review****Syahrir Akil^{1*}, Erni Damayanti², Daud Samsudewa³, Ashariah Hapila⁴, Hikmayani Iskandar⁵**¹Graduate School of Food Technology, Djuanda University, Bogor, Indonesia²Department of Animal Production, Faculty of Animal Science, Universitas Hasanuddin, Makassar, Indonesia³Department of Animal Science, Faculty of Animal and Agricultural Sciences, Universitas Diponegoro, Semarang, Indonesia⁴Graduate School of Animal Nutrition and Feed Science, Faculty of Animal Science, IPB University, Bogor, Indonesia⁵Research Center for Applied Zoology, National Research and Innovation Center Agency (BRIN), Bogor, Indonesia*Email: syahrirakil@unida.ac.id**ABSTRACT**

Zinc is a vital micronutrient involved in various reproductive processes in cattle. This review aims to evaluate the effects of zinc supplementation on cattle reproductive performance by analyzing research articles sourced from Scopus and PubMed databases. Findings indicate that zinc supplementation, provided in either organic or inorganic forms, positively influences reproductive parameters. In bulls, zinc enhances sperm motility, viability, and ejaculate volume. In female cattle, supplementation modulates endocrine hormone levels during both pre-calving and post-calving periods. Moreover, appropriate zinc dosages and supplementation durations contribute to improved reproductive efficiency in cows. These results support the conclusion that zinc supplementation is effective in enhancing reproductive performance in cattle.

Keywords: *Dosage, Reproductive Performance, Cattle, Zinc***ABSTRAK**

Zink merupakan mikronutrien penting yang berperan dalam berbagai proses reproduksi pada sapi. Review ini bertujuan untuk mengevaluasi pengaruh suplementasi zink terhadap performa reproduksi sapi melalui tinjauan artikel penelitian yang diperoleh dari basis data Scopus dan PubMed. Hasil menunjukkan bahwa suplementasi zink, baik dalam bentuk organik maupun anorganik, memberikan efek positif terhadap parameter reproduksi. Pada pejantan, zink meningkatkan motilitas, viabilitas, dan volume ejakulasi sperma. Pada sapi betina, suplementasi zink memodulasi kadar hormon endokrin baik pada masa pra-kelahiran maupun pasca-kelahiran. Selain itu, pemberian zink dengan dosis dan durasi yang tepat dapat meningkatkan efisiensi reproduksi pada sapi betina. Kesimpulannya, suplementasi zink efektif dalam meningkatkan performa reproduksi sapi.

Kata Kunci: *Dosis, Performa Reproduksi, Sapi, Zink*

INTRODUCTION

Zinc (Zn) is an essential trace mineral required for optimal growth, immune competence, and reproductive performance in cattle (Nielsen, 2012). Functioning as a cofactor for more than 200 metalloenzymes, zinc plays a vital role in numerous biological processes, including protein synthesis, antioxidant defense, and the structural stabilization of DNA and RNA (Smith and Akinbamizo, 2000). It is also critical in regulating cell division, differentiation, and nucleic acid metabolism throughout the life cycle (McDonald et al., 2007).

Zinc contributes significantly to hormonal activity, influencing the synthesis, storage, and secretion of key reproductive hormones. Additionally, it supports immune function and maintains electrolyte and acid–base balance, which are essential for maintaining systemic homeostasis (McDonald et al., 2007). Given these roles, even marginal zinc deficiency can disrupt physiological functions and negatively impact reproductive efficiency.

Zinc deficiency has been widely associated with various reproductive impairments, including reduced fertility, prolonged postpartum anestrus, retained placenta, increased services per conception, and early embryonic loss. These reproductive dysfunctions have been reported in multiple species, including cattle, rats, goats, monkeys, poultry, and humans (Nawan et al., 2018; Liu et al., 2019; Galarza et al., 2020; Javari et al., 2021; Vickram et al., 2021). At the cellular level, zinc influences spermatozoa through its regulatory role in capacitation and the acrosome reaction—two critical events required for successful fertilization (Michailov et al., 2014).

In cattle, semen quality is a key determinant of reproductive success, particularly in artificial insemination (AI) programs. Parameters such as motility, morphology, concentration, membrane integrity, and DNA stability are commonly used to assess semen quality. Zinc has been shown to influence many of these parameters, primarily through its antioxidant properties and its involvement in membrane stabilization and chromatin condensation. Despite growing interest in the role of trace minerals in male

fertility, there remains a need to consolidate existing evidence on the effects of zinc supplementation on semen quality in bulls.

Therefore, the objective of this review is thus to describe current research on the effects of zinc supplementation on reproductive performance in cattle. Elucidating the physiological role of zinc, particularly in relation to male reproductive function, is critical for the formulation of precise nutritional strategies aimed at optimizing reproductive efficiency. Such insights are expected to contribute to enhanced fertility outcomes and improved productivity in cattle breeding programs.

DATA COLLECTION

A comprehensive literature search was conducted using Publish or Perish (PoP) software version 8 to identify relevant studies on the effects of zinc supplementation on reproductive performance in cattle. The search was performed using the keywords “*zinc supplementation*” and “*reproductive performance in cattle*” across three major databases: PubMed, ScienceDirect, and Google Scholar. The inclusion criteria encompassed peer-reviewed articles published between 2007 and 2025. All retrieved references were exported in RIS format and processed using the CADIMA web-based systematic review tool (CADIMA, 2020). After removing duplicates, a total of 30 unique records were identified. These records were randomized and subjected to full-text screening based on predefined eligibility criteria. Ultimately, five studies were included in the final review, all of which met the relevance and quality standards within the scope of reproductive performance.

FORMS OF ZINC SUPPLEMENTATION

Zinc supplementation in cattle diets is commonly administered in two primary forms: inorganic and organic. Inorganic sources such as zinc sulfate and zinc oxide have traditionally been used due to their availability and cost-effectiveness (NRC, 2012). However, these forms often exhibit lower bioavailability because of their tendency to form insoluble complexes in the

gastrointestinal tract, which limits absorption. In contrast, organic zinc sources—such as zinc methionine, zinc proteinate, and zinc amino acid chelates—demonstrate higher bioavailability and greater biological efficacy due to their improved stability and absorption across the intestinal mucosa. Comparative studies have shown that

organic zinc is more effective in enhancing reproductive parameters, likely due to its superior capacity to reach target tissues, including the testes and accessory sex glands. The choice of zinc form is thus critical for optimizing its reproductive benefits (Fallah et al., 2018).

Table 1. Zinc supplementation and its effects on reproductive performance in cattle

No	Cattle	Zinc Type	Dose	Reference
1	Crossbred cattle (<i>Bos indicus</i> x <i>Bos taurus</i>)	Inorganic and organic	0, 35, and 70 ppm	Kumar et al., 2006
2	Kankrej bulls	Organic	100 ppm	Patil et al., 2023
3	Frieswal bulls	Inorganic and organic		Chauhan et al., 2021

Zinc supplementation, whether in inorganic or organic form, significantly improves both the quantitative and qualitative characteristics of semen in crossbred bulls. According to Kumar et al. (2006), dietary inclusion of zinc resulted in a statistically significant increase in semen volume, sperm concentration, progressive motility, and viability after six months of supplementation. Furthermore, functional assessments such as the bovine cervical mucus penetration test (BCMPT) and hypo-osmotic swelling test (HOST) revealed enhanced sperm membrane integrity and functional competence in zinc-treated bulls. Enzymatic analysis demonstrated elevated alkaline and acid phosphatase activities in the seminal plasma of supplemented groups, suggesting improved secretory function of the accessory sex glands. Conversely, a notable reduction in GOT and GPT activity in bulls receiving organic zinc (zinc propionate) may indicate reduced oxidative stress and enhanced seminal plasma quality. Additionally, serum testosterone levels were significantly higher in bulls supplemented with higher doses of zinc, particularly in the group receiving organic zinc, highlighting zinc's role in modulating the endocrine axis involved in reproductive function. Collectively, these findings support the beneficial effects of zinc—especially in organic form—in enhancing reproductive efficiency in male cattle.

The supplementation of 100 ppm zinc propionate in Kankrej bulls markedly improved both the quantitative and qualitative parameters of semen, underscoring the

essential role of zinc in male reproductive physiology. Moreover, improvements in mass motility and the percentage of live sperm post-supplementation indicate enhanced sperm vitality and functionality, which are critical determinants of fertility. These findings align with zinc's known role as a cofactor in testicular steroidogenesis and membrane stabilization, particularly in the context of testosterone regulation via Leydig cell function. Given the widespread zinc deficiency in Indian soils, the study emphasizes the practical relevance of targeted mineral supplementation in improving semen quality, thereby supporting reproductive performance and efficiency in indigenous cattle breeds such as Kankrej (Patil et al., 2023).

Chauhan et al. (2021) reported that combined dietary supplementation of selenium (0.20 mg/kg DMI) and zinc (35 mg/kg DMI) in Frieswal bulls significantly enhanced semen quality and sperm output compared to non-supplemented controls. Bulls receiving the combined treatment (T2) exhibited marked improvements in semen volume, sperm concentration per ejaculate, gross motility, and progressive motility, suggesting enhanced spermatogenic efficiency and sperm functional integrity. These parameters are crucial indicators of male fertility potential, and the synergistic effects of selenium and zinc likely support antioxidative defense mechanisms and testosterone biosynthesis, both essential for optimal sperm development and motility. Although sperm concentration per milliliter was unexpectedly highest in the control group (T0),

the total sperm yield per ejaculate and motility metrics were superior in the supplemented groups, indicating that selenium and zinc may contribute more significantly to overall reproductive output than to sperm density alone. These results reinforce the relevance of trace mineral supplementation in improving reproductive performance in breeding bulls, particularly under practical feeding conditions (Chauhan et al., 2021).

EFFECT ZINC ON SEMEN QUALITY PARAMETERS

Zinc plays a pivotal role in maintaining and improving semen quality parameters in bulls. It is a crucial structural and functional

component of numerous enzymes involved in spermatogenesis. Supplementation with zinc has been associated with significant improvements in sperm concentration, motility, viability, and membrane integrity (Esfiohki et al., 2023). Several studies report that zinc supports the maintenance of acrosomal integrity, reduces sperm abnormalities, and limits DNA fragmentation, all of which are essential for fertilization capability (Oliveira et al., 2004; Fadl et al., 2022). Moreover, zinc contributes to the stabilization of sperm chromatin and the protection of membrane lipids from oxidative damage through its role as an antioxidant.

Table 2. Zinc effects on cattle semen quality

Parameter	Effect on Semen Quality	Reference
Sperm Motility	Significant increase in total and progressive motility	Galarza et al., 2020
Sperm Morphology	No significant change; trend towards improvement	Kumar et al., 2006
Sperm Viability	Increased viability and membrane integrity	Kumar et al., 2006; Esfiohki et al., 2023
Sperm Concentration	Increased sperm concentration and ejaculate volume	Kumar et al., 2006
DNA Integrity	Likely protective via antioxidant activity	Esfiohki et al., 2023

HORMONAL REGULATION AND TESTICULAR FUNCTION

Zinc is essential for the regulation of testicular function and endocrine signaling pathways that govern male fertility. It directly influences the synthesis and secretion of testosterone by Leydig cells and supports the function of Sertoli cells, which are responsible for nurturing developing germ cells. Zinc also modulates the secretion of gonadotropins, particularly luteinizing hormone (LH) and follicle-stimulating hormone (FSH), through its involvement in hypothalamic–pituitary axis regulation. Deficiency in zinc has been associated with impaired testicular histology, reduced seminiferous tubule diameter, and decreased sperm output. Conversely, adequate zinc levels promote the proliferation and differentiation of spermatogenic cells, supporting efficient spermatogenesis and optimal testicular morphology.

genic cells, supporting efficient spermatogenesis and optimal testicular morphology.

The mineral zinc (Zn) plays a crucial role in supporting male reproductive function, notably by stimulating Leydig cells to produce testosterone, which in turn induces libido (Fallah et al., 2018). During spermatogenesis, zinc is essential for ribonuclease activity at the early stages of spermatogenesis and contributes to the maturation of spermatozoa throughout the process, ultimately enhancing sperm motility in the later stages. Zinc also exhibits antioxidant properties, functioning to neutralize reactive oxygen species (ROS). Furthermore, zinc protects germ cells from oxidative stress, which can otherwise impair the structure and function of spermatozoa (Sahu et al., 2020).




Figure 1. A model describing the mechanisms by which Zn^{2+} stimulates hyperactivated motility (HAM) and acrosomal exocytosis (AE): Zn^{2+} binds to and activates the GPR39 receptor, which in turn stimulates transmembrane adenylyl cyclase (tmAC) to produce cyclic AMP (cAMP). This increase in cAMP activates the sodium/hydrogen exchanger (NHE), raising intracellular pH (pHi) and triggering CatSper activation. The resulting influx of Ca^{2+} , along with bicarbonate ions (HCO_3^-), activates soluble adenylyl cyclase (sAC), further elevating intracellular cAMP levels. Elevated cAMP activates protein kinase A (PKA), which then initiates the Src-EGFR-PLC signaling cascade, leading to inositol triphosphate (IP_3) production. IP_3 mobilizes Ca^{2+} from intracellular acrosomal stores, thereby amplifying intracellular calcium concentrations and promoting hyperactivation. Concurrently, PKA activation stimulates phospholipase D1 (PLD1), facilitating F-actin polymerization during sperm capacitation. Prior to AE, PLC hydrolyzes phosphatidylinositol 4,5-bisphosphate (PIP_2), leading to the release of the actin-severing protein gelsolin into the cytosol. Activated by Ca^{2+} , gelsolin induces F-actin depolymerization, thereby enabling acrosomal exocytosis. Adapted from Allouche -Fitoussi and Breitbart (2020).

Table 3. Effect of zinc supplementation on endocrine hormones

Hormone	Effect	References
Leptin (ng/ml)	Increases leptin concentration during pre-calving and post-calving	Chandra et al., 2018
Insulin (ng/ml)	Increases insulin concentration during pre-calving and post-calving	Chandra et al., 2018
Growth Hormone (ng/ml)	Increases growth hormone concentration during pre-calving and post-calving	Chandra et al., 2018
IGF-1(nmol/l)	Increases IGF-1 concentration during pre-calving and post-calving	Chandra et al., 2018
Cortisol (ng/ml)	Increases cortisol concentration during pre-calving and post-calving	Chandra et al., 2018
Triiodothyronine (ng/ml)	Increases triiodothyronine concentration during pre-calving and post-calving	Chandra et al., 2018
Thyroxine (ng/ml)	Increases thyroxine concentration during pre-calving and post-calving	Chandra et al., 2018

Zinc supplementation during the periparturient period has a positive effect on female cattle by supporting their condition

before and after calving. It has been shown that zinc reduces stress and helps maintain endocrine hormone concentrations.

Zinc supplementation during the pre-calving and post-calving periods, at appropriate doses and durations, can enhance the immune system (Chandra et al., 2018). Zinc functions as an antioxidant and supports immune defense through its critical role in cell replication and proliferation (Wessels et al., 2017). It also protects cell membranes from inflammatory responses and, mediated by phosphorus, facilitates the migration of blood neutrophils (Finamore et al., 2008). The inclusion of zinc in the diet has been shown to improve immunity in both cows and calves (Dang et al., 2011; Wo et al., 2022).

DOSE AND DURATION OF SUPPLEMENTATION

The efficacy of zinc supplementation on reproductive outcomes is influenced by

both the administered dose and the duration of supplementation. Studies have employed varying concentrations, generally ranging from 30 to 120 mg of zinc per kg of dry matter (DM), with positive reproductive effects often observed at mid-range levels (e.g., 60–80 mg/kg DM). Prolonged supplementation for periods of 8 to 12 weeks tends to yield significant improvements in semen quality, aligning with the full cycle of spermatogenesis in bulls. However, excessively high doses may pose toxicity risks or lead to antagonistic interactions with other trace minerals such as copper. Therefore, precise dosing based on animal requirements, zinc source, and physiological status is essential to maximize reproductive benefits without compromising overall mineral balance.

Table 4. Zink dosage and duration

	Zinc Dosage	Duration (Days)	Effect	References
IVF Medium	0.4, 0.8 and 1.2 (µg/mL)	6	Improved viability, functional sperm membrane integrity (HOST), and sperm motility	Galarza et al., 2020
IVF medium	0.4, 0.8 and 1.2 (µg/mL)	-	Increased fertilization rate	Galarza et al., 2020
IVF medium	0.4, 0.8 and 1.2 (µg/mL)	-	Enhanced cleavage, blastocyst, and hatching rates	Galarza et al., 2020
Pre-calving	80 ppm, 120 ppm	45	Reduced postpartum estrus interval and service period; improved conception rate in dairy cows	Patel et al., 2016
Post-calving	80 ppm, 120 ppm	45	Improved reproductive efficiency in dairy cows	Patel et al., 2016
Beef Cow	400 mg	9	Increased preovulatory follicle size, plasma progesterone concentration, and pregnancy rates	Anchordoquy et al., 2019

Zinc supplementation in IVF medium improves sperm quality in vitro. Zinc acts as an activator of enzymes involved in energy metabolism, such as dehydrogenases and transferases. The activity of these enzymes generates energy required to support sperm motility, ensuring optimal movement during

the fertilization process (Kerns et al., 2018). Additionally, zinc plays a crucial role as a hormonal modulator in the maturation of oocytes and the formation of cumulus-oocyte complexes (COCs) in cattle (Pascua et al., 2020).

CONCLUSION

Zinc supplementation at specific dosages and durations can improve the efficiency of reproductive performance in cattle.

REFERENCES

- Allauche-Fitoussi D and Breitbart H. 2020. The role of zinc in male fertility. *Int J Mol Sci*, 21(20):7796.
- Anchordoquy, J. M., Anchordoquy, J. P., Galarza, E. M., Farnetano, N. A., Giulliodori, M. J., Nikoloff, N., ... & Furnus, C. C. (2019). Parenteral zinc supplementation increases pregnancy rates in beef cows. *Biological Trace Element Research*, 192, 175-182.
- CADIMA. 2020. CADIMA. Jki - julius kühn-institut – Federal research centre for cultivated plants Cadima vers. 2.2 – Dec 2020 copyright(©) 2012-2020. Available at: <https://www.cadima.info/index.php/site/about>.
- Chandra, G., Aggarwal, A., Kumar, M., & Singh, A. K. (2018). Effect of zinc and vitamin E supplementation on hormones and blood biochemicals in peripartum Sahiwal cows. *Journal of Trace Elements in Medicine and Biology*, 50, 489-497
- Chauhan HS, Rajkumar, Kumar S, Kumar A, and Sahu DS. 2021. Inclusion of selenium and zinc supplementation on semen quality of frieswal cattle bulls. *J Exo zOOL iNDIA*, 24(2):1017-1020.
- Dang, A. K., Prasad, S., De, K., Pal, S., Mukherjee, J., Sandeep, I. V. R., ... & Prakash, B. S. (2013). Effect of supplementation of vitamin E, copper and zinc on the in vitro phagocytic activity and lymphocyte proliferation index of peripartum Sahiwal (*Bos indicus*) cows. *Journal of Animal Physiology and Animal Nutrition*, 97(2), 315-321.
- Esfiochi SHM, Norouzian MA, and Najafi A. 2023. Effect of different sources of dietary zinc on sperm quality and oxidative parameters. *Front Vet Sci*, 10:1134244.
- Fadl AM, Abdelnaby EA, and El-Sherbiny HARI. 2022. Supplemental dietary zinc sulphate and folic acid combination improves testicular volume and haemodynamics, testosterone levels, and semen quality in rams under heat stress conditions. *Reprod Domestik Anim*, 57:567-576.
- Fallah A, Mohammad-Hasani A, and Colagar AH. 2018. Zinc is an essential element for male fertility: A review of Zn roles in men's healths, germination, sperm quality, and fertilization. *J Reprod Infertil*, 19(2):69-81.
- Fallah, A., Mohammad-Hasani, A., & Colagar, A. H. (2018). Zinc is an essential element for male fertility: a review of Zn roles in men's health, germination, sperm quality, and fertilization. *Journal of reproduction & infertility*, 19(2), 69.
- Finamore, A., Massimi, M., Devirgiliis, L. C., & Mengheri, E. (2008). Zinc Deficiency Induces Membrane Barrier Damage and Increases Neutrophil Transmigration in Caco-2 Cells^{1, 2}. *The Journal of nutrition*, 138(9), 1664-1670.
- Galarza EM, Lizarraga RM, Anchordoquy JP, Farnetano NA, Furnus CC, Fazzio LE, and Anchordoquy JM. 2020. Zinc supplementation within the reference ranges for zinc status in cattle improves sperm quality without modifying *in vitro* fertilization performance. *Anim Reprod Sci*, 221:106595.
- Jafari M, Irani M, and Rezaeipour V. 2021. Effect of different dietary zinc sources on the semen quality, testicular histology and sex hormone concentration in broiler breeder roosters. *Italian J Anim Sci*, 26,1:489-495.
- Kerns, K., Zigo, M., & Sutovsky, P. (2018). Zinc: A necessary ion for mammalian sperm fertilization competency. *International journal of molecular sciences*, 19(12), 4097.
- Kumar N, Verma RP, Singh LP, Varshney VP, and Dass RS. 2006. Effect of different levels and sources of zinc supplementation on quantitative and qualitative semen attributes and serum testosterone level in crossbred cattle (*Bos indicus* x *Bos taurus*). *Reprod Nutr Dev*, 46(6):663-675.
- Liu H, Sun Y, Zhao J, Dong W, and Yang G. 2019. Effect of zinc supplementation on semen quality, sperm antioxidant ability, and seminal and blood plasma

- mineral profiles in Cashmere goats. *Bio Tra Elem Res*,196:438-445.
- Mcdonald P, Edwards RA, Greenhalgh JFD, Morgan CA, Sinclair LA, and Wilkinson RG. 2007. *Animal Nutrition*. p.130.
- Michailov Y, Ickowicz D, and Breitbart H. 2014. Zn²⁺-stimulation of sperm capacitation and of the acrosome reaction is mediated by EGFR activation. *Dev Biol*, 396(2):246-255
- National Research Council (NRC). 2012. *Nutrient requirements of swine*. 12 ed. National Academy Press, Washington DC, USA.
- Nawab A, Zhao Y, Ibtisham F, Li G, Xiao M, Wu J, Liu W, Tang S, and An L. 2018. The potential effect of zinc deficiency on reproductive profile of male rat and its possible consequences. *Anim Rev*, 5,212-21.
- Nielsen FH. 2012. History of zinc in agriculture. *Adv. Nutr.* 3 (6), 783–789.
- Oliveira C, Badu C, Ferreira W, Kamwa E, and Lana A. 2004. Effects of dietary zinc supplementation on spermatid characteristics of rabbit breeders. In: 8th World Rabbit Congress, Mexico.
- Pascua, A. M., Nikoloff, N., Carranza, A. C., Anchordoquy, J. P., Quintana, S., Barbisán, G., ... & Furnus, C. C. (2020). Reproductive hormones influence zinc homeostasis in the bovine cumulus-oocyte complex: Impact on intracellular zinc concentration and transporters gene expression. *Theriogenology*, 146, 48-57.
- Patel, B., Kumar, N., Jain, V., Ajithakumar, H. M., Kumar, S., Raheja, N., ... & Singh, S. V. (2017). Zinc supplementation improves reproductive performance of Karan-Fries cattle. *Indian Journal of Animal Reproduction*, 38(1), 20-22.
- Patil SS, Rathod BISA, Pawar MM, Chaudhary AB, and Modi CP. 2023. Effects of organic zinc supplementation on semen production and quality in Kankrej bulls. *Asian J Dairy Food Res*, 42(2):183-186.
- Sahu, C., Dwivedi, D. K., & Jena, G. B. (2020). Zinc and selenium combination treatment protected diabetes-induced testicular and epididymal damage in rat. *Human & experimental toxicology*, 39(9), 1235-1256.
- Smith OB and Akinbamizo OO. 2000. Micronutrients and reproduction in farm animals. *Anim Reprod Sci*, 60-61: 549-560.
- Vickram S, Rohini K, Srinivasan S, Veenukumari DN, Archana K, Anbarasu K, Jeyanthi P, Thanigaivel S, Gulothungan G, Rajendiran N, and Srikumar PS. 2021. Role of zinc (Zn) in human reoroduction: A journey from initial spermatogenesis to childbirth. *Int J Mol Sci*, 22;22(4):2188
- Wessels, I., Maywald, M., & Rink, L. (2017). Zinc as a gatekeeper of immune function. *Nutrients*, 9(12), 1286.
- Wo Y, Jin Y, Gao D, Ma F, Ma Z, Liu Z, Chu K and Sun P (2022) Supplementation With Zinc Proteininate Increases the Growth Performance by Reducing the Incidence of Diarrhea and Improving the Immune Function of Dairy Calves During the First Month of Life. *Front. Vet. Sci.* 9:911330.