

**ANALYSIS OF THE PHYTOCHEMICAL POTENTIAL OF *Moringa oleifera* LEAF TEA FORMULATION AND NATURAL SPICES AS AN ALTERNATIVE TO STUNTING PREVENTION****Analisis Potensi Fitokimia Formulasi Teh Daun Kelor (*Moringa oleifera*) dan Rempah-Rempah Alami Sebagai Alternatif untuk Pencegahan Stunting****Angga Dwi Prasetyo^{1*}, Moh. Taufik², Bertina Seviana Putri¹, A'ida Fajrin Najwa²**¹Program Studi Bioteknologi, UIN Raden Mas Said Surakarta.

Jl. Pandawa, Pucangan, Kec. Kartasura, Kabupaten Sukoharjo, Jawa Tengah 57168

²Program Studi Teknologi Pangan, UIN Raden Mas Said Surakarta.

Jl. Pandawa, Pucangan, Kec. Kartasura, Kabupaten Sukoharjo, Jawa Tengah 57168

*E-mail: angga.dwiprasetyo@staff.uinsaid.ac.id**ABSTRACT**

Stunting prevention efforts require a locally focused, nutrient-rich, and easily accessible approach. This study aims to analyze the phytochemical potential of moringa tea (*Moringa oleifera*) and various natural spices such as ginger, turmeric, lemongrass, and galangal as natural alternatives in stunting prevention. The research will be carried out from June to December 2024. The methods used include qualitative phytochemical tests to identify active compounds such as flavonoids, tannins, saponins, alkaloids, and terpenoids/steroids; the formulation of moringa leaf tea and natural spices; and a literature review and meta-analysis of the role of these bioactives in supporting the nutritional status and immune system of children. The results of the analysis showed that moringa tea and spices contain a variety of bioactive compounds with antioxidant, anti-inflammatory, and immunostimulant properties that have the potential to support optimal growth of children. Based on the results of meta-analysis and literature studies, it can be concluded that the phytochemical content of Moringa leaf tea has the potential to prevent stunting in mothers and children. Further research is needed to test the effectiveness of the formula and optimal dosage in clinical applications.

Keywords: *Stunting, Moringa oleifera, Natural Spices, Phytochemicals***ABSTRAK**

Upaya pencegahan stunting membutuhkan pendekatan berbasis pangan lokal yang kaya nutrisi dan mudah diakses. Studi ini bertujuan untuk menganalisis potensi fitokimia teh moringa (*Moringa oleifera*) dan berbagai rempah alami seperti jahe, kunyit, serai, dan lengkuas sebagai alternatif alami dalam pencegahan stunting. Penelitian akan dilakukan dari Juni-Desember 2024. Metode yang digunakan meliputi uji fitokimia kualitatif untuk mengidentifikasi senyawa aktif seperti flavonoid, tanin, saponin, alkaloid, dan terpenoid/steroid, formulasi teh daun moringa dan rempah alami, serta tinjauan literatur dan meta-analisis peran bioaktif ini dalam mendukung status gizi dan sistem imun anak. Hasil analisis menunjukkan bahwa teh moringa dan rempah-rempah mengandung berbagai senyawa bioaktif dengan sifat antioksidan, anti-inflamasi, dan imunostimulan yang berpotensi mendukung pertumbuhan optimal anak. Berdasarkan hasil meta-analisis dan studi literatur, dapat disimpulkan bahwa kandungan fitokimia teh rempah daun kelor berpotensi mencegah stunting pada anak. Diperlukan penelitian lebih lanjut untuk menguji efektivitas formula dan dosis optimal dalam aplikasi klinis.

Kata Kunci: *Stunting, Moringa oleifera, Rempah Alami, Fitokimia*

INTRODUCTION

Stunting is one of the global nutrition problems that is being handled by various countries and organizations in the world. Data from the UNICEF/WHO/WB Joint Child Malnutrition Estimates (JME) that in 2021 there were 149 million children under five declared stunted, 45 million children under five wasting, and 39 million overweight toddlers. The Ambitious World Health Assembly targets a 40% reduction in stunting globally across the country. Indonesia is one of the countries with the second highest prevalence of stunting (after Cambodia) in the Southeast Asian region and is ranked 108th out of 132 countries in the world. The results of the 2021 Indonesian Nutrition Status Study state that the prevalence of stunting in toddlers is still high, at 24.4% (Saepudin, 2024).

The incidence of stunting is caused by various factors, one of which is the lack of nutritional intake that occurs in the first 1000 days of life (1000 HPK). Children have the potential to experience stunting since pregnancy due to malnutrition during pregnancy. The cause of stunting also occurs if they suffer from anemia during pregnancy (Khotimah, 2023). Pregnant women need to consume highly nutritious food, both macro and micro, as well as antioxidants. One of the preventive aspects that can be done and become a priority in accordance with the direction of the Ministry of Health is the food aspect.

According to Marhaeni (2021), moringa leaves can be used as food ingredients that have a fairly high nutritional content or the term is superfood. Consuming this superfood can improve a person's nutritional status and immunity because of the many bioactive compounds found in moringa leaves. The results of Minantyo et al. (2019) research show that moringa leaves have a large nutritional content. The protein content of moringa leaves is 26.2 g, calcium is 2,095 mg, iron is 27.1 mg, and provitamin A is quite high, which is 16800 mg (Su & Chen, 2020). In addition, the results of research by More et al. (2020) show that moringa leaves have 40 natural antioxidants. Given the content of these nutrients,

moringa leaves can be used as herbal or traditional medicine (Marhaeni, 2021).

During the period of improving the nutrition of a child affected by stunting, moringa leaves can be used as nutritional support and immune boosters for the body of stunted children so that the health of stunted children can be maintained and avoid disease (Putra et al., 2021). Several researchers who have used moringa leaves as a stunting prevention product, namely Istiqomah and Jannah (2021) make moringa pudding; Ishmah et al. (2023) make moringa nuggets; Kustiani et al. (2022) make moringa flour; and Swari et al. (2023) make moringa ice cream. In addition, the content of natural spices such as ginger (Leslie & Gunawan, 2023), lemongrass (Najmah et al., 2023), galangal (Badriyah et al., 2023) and turmeric (Zakaria et al., 2023) also have phytochemical compounds that have potential in the process of preventing stunting disease.

Based on several previous studies (Basri et al., 2021), the use of *Moringa oleifera* combined with natural spices has the potential to be a supplement in pregnant women and toddlers as a prevention of stunting disease. This study aimed to evaluate the potential of moringa leaf tea combined with natural spices as an alternative approach to preventing stunting, based on phytochemical analysis, and to develop a formulation of moringa tea and natural spices as a herbal beverage.

MATERIALS AND METHODS

Optimization and Manufacturing of Natural Spice Moringa Leaf *Simplicia* Powder

The *simplicia* powder begins with the process of washing all ingredients using water with a basin container, then thinly slice the ingredients using a knife on a placemat except moringa leaves, and finally the drying process. Optimization is carried out on moringa leaves to see the best results in terms of color production, then the results of optimization of the drying method that have been selected are used for natural spices, method optimization includes drying using sunlight for 2 days, using aluminum foil base, using an oven with a temperature of

40°C for 16 hours with an aluminum foil base and using a dehydrator with a temperature of 55°C for 5 hours with a cloth gauze base. After the material is dry, then the material is smoothed into powder using a grinder at a

different time, the finished powder is stored in ziplock plastic, then sealed so that it is durable in storage. The process of making spiced Moringa tea powder is shown in **Figure 1**.




Figure 1. Process Optimization and Manufacturing of Natural Spice Moringa Leaf Simplicia Powder

Moringa Leaf Tea Formulation and Natural Spices

The powder that has been dried, then formulated by calculating the composition of each ingredient, taking into account simple organoleptics, including the color, taste and aroma of spiced moringa leaf tea. The ingredients include moringa powder, ginger powder, lemongrass powder, turmeric powder, galangal powder, and sugar (Table 1). The results of the formulation were then tested in simple organoleptics by packaging them into tea bags and then brewed in hot water, choosing which formulation met the characteristic criteria of the formula, then the formulation results were packaged into tea bags of tea bags of moringa leaves and natural spices.

Moringa Leaf Extraction and Natural Spices

Moringa leaf powder and natural spices are taken as much as 100 g and put into the erlenmeyer to be macerated and

dissolved with 500 mL of 96% ethanol in a ratio of 1:5. Erlenmeyer is covered with aluminum foil and wrapped in plastic wrap, then put into an incubator shaker at 37 °C for 24 hours and a speed of 160 rpm. The sample was filtered after 24 hours and the first filtrate and pulp were obtained, then the first pulp was macerated with 96% ethanol for 24 hours. The sample was filtered with filter paper and a second filtrate and pulp were produced (Wijanarko et al., 2020).

The filtrate obtained from maceration is fed into a rotary evaporator at a temperature of 60 °C and a speed of 65 rpm. The viscous extract is put into a porcelain cup and covered with aluminum foil, then ovened at a temperature of 50 °C and observed for 24 hours until the result is obtained in the form of a paste (Wijanarko et al., 2020). The simplicia rendition is calculated by the formula (Wijanarko et al., 2020):

$$\text{Extract weight} = (\text{cup weight} + \text{extract}) - \text{empty cup weight}$$

Phytochemical Analysis of Moringa Leaf Tea and Natural Spices

Sample preparation for qualitative phytochemical analysis including flavonoids, tannins, saponins, and alkaloids was carried out by weighing 50 mg of bandotan leaf extract and then dissolved with 5 mL of 96% ethanol so that the concentration was obtained 10,000 ppm (Ikalinus, et al., 2015). Flavonoid analysis was carried out by diluting the sample to a concentration of 4000 ppm, then taking as much as 2 mL and adding 3 mL of aquaade. Then it is brought to a boil and added with 15 mg of Mg powder, 2 drops of HCl 5N, and 10 drops of amyl alcohol. Flavonoid positive results are yellow, red, or orange in the amyl alcohol layer (Syafitri et al., 2014).

Alkaloid analysis was done by taking 2 mL of extract solution with a concentration of 10,000 ppm and supplemented with 3 mL of aquaade. The solution was added with 1 mL of chloroform, 2 mL of ammonia and then beaten and added with 5 mL of HCl 2 N. The solution was put into 3 different test tubes and each test tube was given 2-3 drops of Dragendrof, Mayer, and Wagner reagents. The positive results of the Dragendrof reaction were orange deposits, the Mayer reaction formed white deposits, and the Wagner reaction formed brown to yellow deposits (Januarti et al., 2017).

Tannin analysis was carried out with the extract diluted to 4000 ppm. The solution is taken as much as 1 mL then added to 5 mL of aqueducts, then boiled and added with 1 % FeCl₃ as many as 5 drops. Positive results were formed in greenish-brown or blackish-blue color (Januarti et al., 2017). Saponin analysis was carried out by extracting a concentration of 4000 ppm as much as 2 mL, plus 10 mL of aqueduct and whipped

for 30 seconds. The positive result was that foam was formed with a height of about ± 1 cm which lasted for approximately 30 seconds (Januarti et al., 2017).

Triterpenoid and steroid analysis was performed by means of diluted extracts up to 4000 ppm. The test is carried out by taking a test solution with a dropper and dripping it in two different drop plates. The first hole is for control and the second hole is for the test solution. Added a drop of anhydrous acetate and 1 drop of H₂SO₄ on the sample. Positive results of steroid analysis are indicated in green, while triterpenoid tests are red or purple (Ikalinus et al., 2015).

Data Analysis of Phytochemical Content for the Prevention of Stunting

The data from the results of phytochemical analysis were qualitatively analyzed and the study literature was then carried out to see the relationship and relationship in the prevention of stunting disease. Furthermore, a description analysis was carried out to see the potential of natural spiced moringa tea as a stunting prevention by looking at the relationship of bioactive compounds in natural spiced moringa tea to stunting prevention with metaanalysis and literature review.

RESULT AND DISCUSSION

Optimization and Manufacturing of Natural Spice Moringa Leaf Simplicia Powder

Based on the results of optimizing drying moringa leaves using 3 different methods, namely drying using dehydrators, ovens and sunlight, after being pureed, different results were obtained in terms of color. The results of the optimization of moringa leaf powder are presented in **Figure 2**.




Figure 2. Moringa Leaf Powder using Dehydrator (Left), Oven (Middle), and Sunlight (Right)

Based on the optimization results, the dehydrator drying method was chosen as the method of making natural spice powder, then the powder that has been made is

packed using ziplock plastic and sealed so that the simplicia powder is durable. The results of making powder using dehydrators on natural spices are presented in **Figure 3**.



Figure 3. Ginger Powder (Left), Turmeric, Galangal, Lemongrass and Moringa Leaves (Right)

The results of simplicia powder are then carried out a formulation process to obtain the right composition of spiced moringa tea based on the results of the formula's characteristics. Based on figure 2, the results of the observation of moringa leaf powder from the three methods can be seen that moringa leaf powder dried using a dehydrator has a greener color compared to using an oven and sunlight. Crownful (2024), stated that the drying method using a dehydrator can maintain the color of the product better than other drying methods. This happens because the dehydrator allows for

more precise temperature and humidity control, which reduces chemical reactions that can fade the color of the material. So that for drying natural spices, a dehydrator is used (figure 3).

Moringa Leaf Tea Formulation and Natural Spices

Based on the results of the formulation by paying attention to the aroma, taste and color of each simplicia, 2 formulations of moringa leaf tea and natural spices were obtained with the details of the formula presented in **Table 1**.

Table 1. Moringa Leaf Tea Formulation and Natural Spices

Simplisia Ingredients	Formula 1 (gr)	Formula 2 (gr)
Ginger	1.0	1.0
Tumeric	0.1	0.1
Lemongrass	0.1	0.1
Galangal	0.1	0.1
Sugar	7.7	7.7
Moringa	1.0	0.75
Total	10	10

Based on the formula that has been obtained, then the characteristics of the formula are carried out with four panelists

covering aspects of color, taste and aroma of the formula. The results of the characteristic formula are presented in **Table 2**.

Table 2. Sensory Characteristics of Moringa Leaf Tea Formula and Natural Spices

Formula	Organoleptic Test Aspects	Organoleptic test results
Formula 1	Color	Deep green
	Flavor	Bitter
	Scent	Pungent spices
Formula 2	Color	Bright green
	Flavor	Slightly bitter
	Scent	Slightly Pungent spices

The formulation of moringa leaf powder (*Moringa oleifera*) requires special attention in maintaining the taste, nutritional content, and solubility of the powder. In this process, additives such as lemon, ginger, or honey are often used to improve the taste and add to the health benefits of the product (Nurfina & Sutji, 2021). In this study, several ingredients were used to improve the taste and also improve its functional properties, namely ginger, turmeric, galangal and lemongrass. Sensory tests are essential in determining functional beverage formulas,

as sensory evaluations such as taste, aroma, color, and texture help ensure the product is accepted by consumers and has good organoleptic quality (Putri et al., 2019; Wibowo & Hidayat, 2020). There are two formulas that are tested by the sensors in this service, namely formula 1 and formula 2 (Table 2). Formula 2 has a larger proportion of moringa leaf powder. The results of sensory tests show that formula 2 has better sensory characteristics compared to formula 1.

Extraction and Phytochemical Analysis of Natural Spiced Moringa Leaf Tea Rendemen

Table 3. Composition of sample Moringa leaf spice tea extract

No	Sample	Powder Weight (g)	Extract Weight (g)	Yield (%)
1	Moringa Leaf	300	28.43	9.24
2	Turmeric	300	30.43	10.37
3	Ginger	300	29.74	10.57
4	Galangal	300	29.98	9.67
5	Lemongrass	300	28.64	9.37

Qualitative and Quantitative Phytochemical Analysis

The results of qualitative and quantitative phytochemical screening of Moringa

leaf and nature spices extracts can be seen in **Table 4** and **Table 5**.

Table 4. Quantitative phytochemical analyses of Moringa leaf and nature spices extracts.

Test	Method	Test Result
Alkaloid	Meyer	+
	Wagner	+
	Dragendroff	+
Tannin	Blackish-blue color	+
Saponin	Foam formed 1 cm	+
Total Flavonoid	Yellow color forms on the amyl alcohol layer	+
Triterpenoid or steroids	Formed a bluish green color	+

Note: (+): positive, (–): negative.

Table 5. Quantitative phytochemical analyses of Moringa leaf and nature spices extracts

Test	Method	Result
Flavonoid	Spektrofotometer UV-Vis	0,78% b/b
Fenol	Spektrofotometer UV-Vis	35,99% b/b
Saponin	Spektrofotometer UV-Vis	2,88% b/b
Tannin	Spektrofotometer UV-Vis	0,91% b/b
Alkaloid	Spektrofotometer UV-Vis	0,24% b/b
Curcuminoid	Spektrofotometer UV-Vis	0,50% b/b

Alkaloid analysis showed positive results, namely in Meyer reagent there was white precipitation and in Dragendroff reagent there was orange precipitation, while in Wagner reagent brown precipitation was found. The results of analysis tanin showed positive results of tannins with the addition of FeCl_3 , Fe^{3+} ions with O- atoms forming covalent bonds because in the OH- function group released the H atom and there was a change in color. The results of positive saponin analysis were characterized by the formation of foam as high as ± 1 cm. The reaction that occurs is that saponins will bind to water to form hydrophilic groups and hydrophobic groups will bond with each other with air so that they can form foam (Oluwafemi et al., 2022).

The results of the analysis showed that extract was positive on the flavonoid test which was characterized by the formation of yellow color on the amyl alcohol layer. The results of the positive analysis of ethanol extract are the formation of a bluish-green color. The change in color of the extract to green is due to the formation of conjugated double bonds on steroids or brown and orange or purple colors on triterpenoids (Ahmed et al., 2022).

The use of ethanol solvents in maceration uses 96% ethanol, harmless safe properties, and non-toxic (Azis et al., 2014). Ethanol solvents are also extracting compounds with low molecular weight such as saponins and flavonoids and the concentration of 96% ethanol in herbal medicine preparations has the highest percentage of yield (Arifianti et al., 2014). The diffusion that occurs in the maceration process occurs in the concentration of the solution in the cell and the solution outside the cell, so that the high concentration solution in the cell will come out and be replaced by a low concentration solution so that the balance occurs inside the cell and outside the cell. The extraction time also affects because the longer the maceration, the higher the yield produced. The longer the reaction between the sample and the solvent used, the better the penetration of the solvent entering the sample so that the more compounds that

diffuse out of the cell will be (Wijaya et al., 2018; Hartoyo, et al., 2025).

Based on the results of extraction and phytochemical qualitative tests in table 3, it was found that the phytochemical content in the composition of natural spiced moringa tea (lemongrass, ginger, galangal and turmeric) was positive for saponins, tannins, flavonoids, alkaloids and triterpenoids as well as steroids. This is shown by the positive results in the phytochemical qualitative test shown in table 3. Furthermore, based on the qualitative results of phytochemicals, an analysis of their potential for disease prevention was carried out through metaanalysis and literature studies. Saponins have the potential to interfere with nutrient absorption and cause large amounts of digestive problems. However, at low levels of saponins can reduce the risk of nutrient malabsorption and indigestion (Zhang et al., 2023). Alkaloids have medicinal properties but can be toxic in high amounts. Lower alkaloid concentrations can reduce the potential for toxic effects (Jegade et al., 2024). Alkaloids help to raise immune function (immunomodulators), which is especially important for malnourished individuals (Oluwafemi et al., 2022).

Flavonoids, have antioxidant properties and are essential in protecting against oxidative stress, which is often elevated in malnourished individuals. Antioxidants can also be used in warding off free radicals that may be produced during malnutrition. Antioxidants can help reduce inflammation and cell damage (Ahmed et al., 2022). Flavonoids are most abundant in all parts of the plant, one of which is in the leaves and belongs to the polyphenol group. Flavonoids are able to inhibit lipid oxidation and ward off free radicals. Flavonoids have the ability to be antioxidant, anti-inflammatory, antibacterial and anticancer (Najmah, et al., 2023). Tannins, while potentially beneficial as antioxidants, can interfere with protein digestion and iron absorption. Lower tannin levels, can improve protein digestion and iron absorption in malnourished individuals, which is important for recovery (Zhang et al., 2019). The results of the study, Hidayah et al. (2023), show that triterpenoids have significant pharmacological activity, such as

antiviral, antibacterial, anti-inflammatory effects, which act as inhibitors of cholesterol synthesis and as anticancer agents. Cancer is a disease caused by abnormal cells in the body's tissues that grow and develop rapidly and uncontrollably.

Analysis of Phytochemical Content Data for the Prevention of Stunting Disease

Based on the meta-analysis and study literature, several phytochemical potentials of natural spiced moringa tea that play a role in the prevention of stunting disease, the results of the phytochemical potential of natural moringa tea are presented in **table 5**.

Table 5. Analysis of Phytochemical Content Data in the relation to preventing stunting

Simplicia	Potential of Phytochemicals to Prevent Stunting	Reference Source
Moringa Leaves	Anti-inflammatory, Anticarcinogenic, immunomodulator, antioxidant, antiproliferation and antiviral	Saini et al., 2014; Shija et al., 2019; Basri et al., 2021; Brar et al., 2022; Afriza et al., 2023; Budiono & Has, 2025
Ginger	Anti-inflammatory, antioxidant, antiproliferative and immunomodulatory	Ali et al., 2008; Kiyama, 2020; Sharma et al., 2021; Unuofin et al., 2021; Fatma & Siddique, 2023; Singh et al., 2023; Edo et al., 2025
Lemongrass	Antioxidant, anticancer, antiproliferation, antimicrobial, antiviral	Gaba et al., 2020; Aboagye et al., 2021; Pan et al., 2022; Adhikary et al., 2024; Alharbi et al., 2024; El-Shan, et al., 2024
Galangal	Antimicrobial, antioxidant, immunomodulator	Das et al., 2020; Sianipar et al., 2021; Alijobair et al., 2022; Ramanunny et al., 2022; Verma & Sharma, 2022; Amalraj et al., 2023
Tumeric	Anti-inflammatory, antioxidant, and immunomodulatory	Permatananda et al., 2021; Abubakar et al., 2022; Oluwafemi et al., 2022; Anyaoku et al., 2023; Cyril, 2023; Hardia et al., 2025

Based on table 5, the potential of moringa leaf tea and spices in the prevention of stunting disease includes Anti-inflammatory, Anticarcinogenic, immunomodulator, antioxidant, antiproliferation and antiviral. One of the important factors that contribute to stunting is chronic inflammation, both derived from recurrent infections and inflammatory bowel conditions such as Environmental Enteric Dysfunction (EED). EEDs cause damage to the intestinal mucosa, interfere with nutrient absorption, and increase intestinal permeability (Santoso et al., 2024). Prolonged systemic inflammation causes linear growth retardation, as it affects growth hormone and bone metabolism. Based on the research in table 4, the phytochemical compounds in spiced moringa tea have a function in inhibiting the production of pro-inflammatory cytokines such as TNF- α , IL-6, and IL-1 β , inhibiting the

activity of COX-2 and prostaglandin enzymes and reducing oxidative stress that triggers inflammation (Sianipar et al., 2021). So that the anti-inflammatory ability of the phytochemical content of spiced moringa tea can reduce systemic inflammation that may interfere with growth and improve intestinal function and nutrient absorption, especially if EED is the cause of stunting (Raczyńska et al., 2025).

In people with stunting and during pregnancy, the immune system will decline in line with the diet and nutrient consumption of the mother and child, this can increase stunting (Kustiani et al., 2022). Moringa leaf tea spice has secondary metabolite compounds with immuno-modulatory activity such as alkaloids, flavonoids, glycosides, polysaccharides, polyphenols, tannins, terpenoids, saponins and sterols (Sianipar, et al., 2021) Types of immunomodulators

consist of three categories: immuno-adjuvant, im-munostimulant, and immunosuppressant. Spiced moringa tea has a role as an im-munostimulant, which can increase the body's resistance to infections, can be worked by the stimulated nonspecific im-mune system (Oluwafemi et al., 2022).

Stunting is the result of chronic disruption of linear growth, one of which is caused by oxidative stress due to persistent inflammation. Oxidative stress occurs when the amount of free radicals (ROS) in the body exceeds the antioxidant capacity (Hasan et al., 2023). This damages intestinal cells, the immune system, and other tissues that are important for growth. According to Zakaria et al (2023) Moringa leaf tea has antioxidant content, this is in accordance with table 4, that flavonoids, tannins and alkaloids are included in the antioxidant group. Antioxidants work in neutralizing free radicals, protecting gut cells and body tissues from damage, reducing inflammation related to oxidative stress, and supporting the immune system, making infections less frequent (Kiyama, 2020; Sharma, et al., 2021). Based on Marhaeni's (2021) re-search, spiced moringa tea contains phyto-chemical compounds that are able to work in the prevention of stunting disease, through maintaining the integrity of intestinal cells, so that nutrient absorption remains optimal, reducing the damaging effects of oxidative stress due to infection or malnutrition, improving nutritional status through the content of micronutrients that support growth, reducing the risk of hormonal disorders that occur due to chronic inflammation and tissue damage (Leslie & Gunawan, 2023).

Based on research conducted by Brar et al. (2022); Afriza et al. (2023); Budiono & Has (2025), shows that the phytochemical compounds contained in spiced moringa tea have antimicrobial, antiviral, anti-inflammatory and anticancer effects that prevent infectious diseases and serious diseases in mothers and children so that they have the potential to cause children to experience stunting. In addition, moringa leaves in spiced moringa tea itself contain many minerals such as calcium, potassium, zinc, magnesium, iron and copper. Vitamins such as beta carotene vitamin A, B vitamins such as folic acid, pyridoxin and nicotinic acid,

vitamins C, D, E are also present in moringa leaves. Moringa leaves also contain low calories so they can be used for diets for people with obesity. Moringa leaves contain many minerals that are essential for growth and development, such as calcium. When compared to milk, 8 ounces of milk contain 300-400 mg of calcium while 1000 mg moringa leaves contain more than 4000 mg of calcium (Kustiani et al., 2022).

The health benefits of moringa leaves are influenced by the nutrients contained in them. In addition to antioxidants, moringa leaves also contain vitamins and minerals, including Vitamin B6, Vitamin B2, Vitamin C, Vitamin A, iron and magnesium. A bowl of moringa leaves ($\pm$ 21 grams) contains 2 grams of vegetable protein (Chalik, 2023). Other studies that have been conducted have also proven that 100 grams of dried moringa leaves contain twice as much protein as yogurt, contain vitamin A seven times higher than carrots, have three times higher potassium content than bananas, have four times higher calcium content than milk, and have seven times higher vitamin C content than oranges. Moringa leaves also contain vitamin B6, iron, magnesium, and riboflavin B2. The diverse content of moringa leaves is what makes moringa leaves get the nickname as a superfood ingredient, which is a functional food that has high nutritional value and is rich in antioxidants that are beneficial to the body, including increasing the fulfillment of nutritional needs in the body. So that spiced moringa tea has the potential to prevent stunting.

CONCLUSION

Moringa tea has the potential as a stunting prevention agent because it contains phytochemicals in the form of alkaloids, tannins, saponins, flavonoids and triterpenoids or steroids that have roles as anti-inflammatory, immunomodulator, antioxidant, antimicrobial, antiviral and anticancer, so that it can prevent stunting in children.

ACKNOWLEDGMENT

Gratitude is addressed to UIN Raden Mas Said Surakarta for providing funding for

research activities through the 2024 LITAPDIMAS KEMENAG RI Grant program.

REFERENCES

- Aboagye, G., Tuah, B., Bansah, E., Tettey, C., & Hunkpe, G. (2021). Comparative evaluation of antioxidant properties of lemongrass and other tea brands. *Scientific African*, 11, e00718. <https://doi.org/10.1016/j.sciaf.2021.e00718>
- Abubakar, I., Danyaya, J. A., Abdullahi, Z., Zubairu, A., Sahabi, A. U., & Ahmad, F. (2022). Phytochemical screening, nutritional and anti-nutritional composition of aqueous rhizome extract of *Curcuma longa*. *J Biotechn Biochem*, 8(2), 1-9. DOI: 10.9790/264X-08020109
- Adhikary, K., Banerjee, P., Barman, S., Bandyopadhyay, B., & Bagchi, D. (2024). Nutritional aspects, chemistry profile, extraction techniques of lemongrass essential oil and it's physiological benefits. *Journal of the american nutrition association*, 43(2), 183-200. <https://doi.org/10.1080/27697061.2023.2245435>
- Afriza, Z. N., Muhani, N., Sari, N., & Sari, F. E. (2023). Use of moringa leaves to meet toddler nutrition in the prevention of stunting. *Jurnal Penelitian Pendidikan Ipa*, 9(12), 11333-11341. <https://doi.org/10.29303/jppipa.v9i12.4325>
- Ahmed, M. H., Vasas, D., Hassan, A., & Molnár, J. (2022). The impact of functional food in prevention of malnutrition. *PharmaNutrition*, 19, 100288. <https://doi.org/10.1016/j.phanu.2022.100288>
- Alharbi, Y. M., Elzahar, K. M., Qahl, S. H., Alghamdi, A. M., Alharbi, A. A., Alanazi, Y. F., & El-Saadony, M. T. (2024). Nutritional significance, antimicrobial, antioxidants, anticancer, and antiviral activities of lemongrass leaves extract and its application as hepatoprotective agent against CCl₄-Induced hepatic injury in rats. *Anais da Academia Brasileira de Ciências*, 96(3), e20230646. <https://doi.org/10.1590/0001-3765202420230646>
- Ali, B. H., Blunden, G., Tanira, M. O., & Nemmar, A. (2008). Some phytochemical, pharmacological and toxicological properties of ginger (*Zingiber officinale* Roscoe): a review of recent research. *Food and chemical Toxicology*, 46(2), 409-420. <https://doi.org/10.1016/j.fct.2007.09.085>
- Amalraj, A., AC, K. V., Jude, S., & Kuttappan, S. (2023). Galangal—Roles in nutraceuticals and functional foods. In *Herbs, Spices and Their Roles in Nutraceuticals and Functional Foods* (pp. 95-109). Academic Press. <https://doi.org/10.1016/B978-0-323-90794-1.00016-8>
- Arifianti L, Oktarina RD, Kusumawati I. 2014. Pengaruh jenis pelarut pengekstraksi terhadap kadar sinensetin dalam ekstrak daun *Orthosiphon stamineus* Benth. *E J Planta Husada*. 2 (1): 1-4.
- Azis T, Febrizky S, Mario AD. 2014. Pengaruh jenis pelarut terhadap persen yield alkaloid dari daun salam india *Murraya koenigii*. *T Kimia*. 2(20):1-6.
- Badriyah, lailatul., Ifandi S., Alfiza I.S. (2023). Analisis kualitatif fitokimia pada rimpang lengkuas putih (*Alpinia galanga* L.) se-bagai antibakteri *Klebsiella pneumo-nia*. *Journal of Herbal, Clinical and Pharmaceutical Science (HER-CLIPS)*, 4(02), 11-17. <http://dx.doi.org/10.30587/her-clips.v4i02.5356>
- Basri, H., Hadju, V., Zulkifli, A., Syam, A., & Indriasari, R. (2021). Effect of *Moringa oleifera* supplementation during pregnancy on the prevention of stunted growth in children between the ages of 36 to 42 months. *Journal of Public Health Research*, 10(2), jphr-2021. <https://doi.org/10.4081/jphr.2021.2207>
- Brar, S., Haugh, C., Robertson, N., Owuor, P. M., Waterman, C., Fuchs III, G. J., & Attia, S. L. (2022). The impact of *Moringa oleifera* leaf supplementation on human and animal nutrition, growth, and milk production: A systematic review. *Phytotherapy Research*, 36(4), 1600-1615. <https://doi.org/10.1002/ptr.7415>

- Budiono, N. D. P., & Has, D. F. S. (2025). The effectiveness of moringa leaves extract to prevent stunting at toddlers aged 23-59 months. *Science Midwifery*, 13(1), 67-73. <https://doi.org/10.35335/mid-wifery.v13i1.1872>
- Chalik, I. (2023). Daun Kelor Miliki Kandungan Protein Berfungsi Cegah Stunting. 6–9. <https://bengkulu.bkkbn.go.id/daun-kelor-miliki-kandungan-protein-berfungsi-cegah-stunting/#:~:text=Dari daunnya yang kecil terdapat,otak anak akibat kekurangan gizi>
- Crownful. (2024). Common Mistakes to Avoid When Using a Food Dehydrator. Diakses dari <https://www.crownful.com>.
- Das, G., Patra, J. K., Gonçalves, S., Romano, A., Gutierrez-Grijalva, E. P., Heredia, J. B., & Shin, H. S. (2020). Galangal, the multipotent super spices: A comprehensive review. *Trends in Food Science & Technology*, 101, 50-62. <https://doi.org/10.1016/j.tifs.2020.04.032>
- Edo, G. I., Igbuku, U. A., Makia, R. S., Isoje, E. F., Gaaz, T. S., Yousif, E., & Umar, H. (2025). Phytochemical profile, therapeutic potentials, nutritional composition, and food applications of ginger: a comprehensive review. *Discover Food*, 5(1), 1-32. <https://doi.org/10.1007/s44187-025-00280-2>
- Fatma, H., & Siddique, H. R. (2023). Research and patents status of selected phytochemicals against cancer: how close and how far?. *Recent Patents on Anti-Cancer Drug Discovery*, 18(4), 428-447. <https://doi.org/10.2174/1574892818666221107113648>
- Gaba, J., Bhardwaj, G., & Sharma, A. (2020). Lemongrass. In *Antioxidants in vegetables and nuts-properties and health benefits* (pp. 75-103). Singapore: Springer Singapore.
- Hardia, L., Akrom, A., Hidayati, T., & Sulistyani, N. (2025). Evaluation of Herbal Honey with Black Cumin and Curcuma xanthorrhiza as an Antioxidant Supplement for Stunting Prevention. *Journal of Public Health and Pharmacy*, 5(1), 42-53. <https://doi.org/10.56338/jphhp.v5i1.5549>
- Hartoyo, V. M., Sidharta, B., & Widiastuti, S. S. (2025). AKTIVITAS ANTIBAKTERI EKSTRAK DAUN BANDOTAN (*Ageratum conyzoides*) TERHADAP *Bacillus cereus* DAN *Salmonella typhi*. *Berita Biologi*, 24(1), 37-49. https://doi.org/10.55981/berita_biologi.2025.5760
- Hasan, T., Ida, N., & Qifni, S. F. (2023). Skrining Fitokimia Dan Uji Aktivitas Antioksidan Ekstrak Etanol Rimpang Kunyit Hitam (*Curcuma Caesia* Roxb.) Asal Luwu Utara Dengan Metode Dpph. *Jurnal Riset Kefarmasian Indonesia*, 5(3), 439-457. <https://doi.org/10.33759/jrki.v5i3.397>
- Hidayah, H., Fatmawati, F., Khairunnisa, J., & Putri, M. H. (2023). Aktivitas triterpenoid sebagai senyawa antikanker. *In-novative: Journal Of Social Science Research*, 3(2), 10168-10183.
- Ikalinus, R., Widyastuti, S. K. dan Setiasih, N. L. E. 2015. Skrining fitokimia ekstrak etanol kulit batang kelor (*Moringa oleifera*). *Indonesia Medicus Veterinus*, 4(1), pp.71-79
- Ishmah, L. Z., Indrasati, A., Rosita, Y. S., Kausar, I., Zurly, S., Nopianti, A., Cahyaninghati, N. W. P., Islamy, M. I., Rosid, A., & Hidayat, N. F. Sosialisasi Stunting dan embuatan Nugget Kelor Sebagai Salah Satu Upaya Pencegahan Stunting. *Jurnal Pengabdian Magister Pendidikan IPA*, 6(3), 626–630. <https://doi.org/10.29303/jpmpi.v6i3.5022>
- Istiqomah, & Jannah, N. L. (2021). Pemanfaatan Daun Kelor Menjadi Sebuah Produk Bernilai Jual di Desa Kesemen Ngoro Mojokerto. *Nasional Seminar for Re-search Community Development*, 5(1), 100–108; 2021
- Januarti, I. B., Santoso, A. dan Razak, A. S. 2017. Ekstraksi senyawa flavonoid daun jati (*Tectona grandis* L.) dengan metode ultrasonik. *Media Farmasi Indonesia*, 12(2), pp.1181-1266.
- Jegede, R. E., Ayeni, G., Abaniwo, R. M., Audu, G. A., & Haruna, A. (2024). Biochemical and Phytochemical Efficacy of Defatted *Moringa oleifera* Seeds in Alleviating Protein-Energy Malnutrition. *bioRxiv*, 2024-10.

- <https://doi.org/10.1101/2024.10.14.618128>
- Khotimah, S. H. H. (2023). Hubungan Pengetahuan Ibu Yang Memiliki Balita Tentang Gerakan 1000 Hari Pertama Kehidupan (HPK) Dengan Kejadian Stunting Pada Balita Di Posyandu Pakuhaji Kabupaten Tangerang: (The Relationship Between Mothers' Knowledge of Toddlers About the First 1000 Days of Life (HPK) Movement and Stunting Incidents in Toddlers at Posyandu Pakuhaji, Tangerang Regency. Indonesian Scholar Journal of Nursing and Midwifery Science (IS-JNMS), 3(02), 1115-1122. <https://doi.org/10.54402/is-jnms.v3i02.399>
- Kiyama, R. (2020). Nutritional implications of ginger: chemistry, biological activities and signaling pathways. The Journal of nutritional biochemistry, 86, 108486. <https://doi.org/10.1016/j.jnutbio.2020.108486>
- Kustiani, A., Adya, A., Nurdin, S. U., & Indriani, Y. (2022). Potensi Tepung Daun Kelor untuk Pencegahan Stunting Sejak Dini.
- Leslie, A. G. J., & Gunawan, S. (2023). Ekstrak jahe merah (*Zingiber officinale* Var. *Rubrum*): uji fitokimia, analisa sidik jari, kapasitas total antioksidan, dan penentuan kadar fenolik. Jurnal Kesehatan Tam-busai, 4(2), 2007-2016.
- Marhaeni, L. S. (2021). Daun Kelor (*Moringa Oleifera*) Sebagai Sumber Pangan Fungsional Dan Antioksidan Lulus. Agrisia, 13(2), 40–53.
- Minantyo, H., Purnomo, H., Winarno, P. S., & Kartikawati, M. (2019). The improvement of nutrition quality and organoleptic characteristics of Indonesian milkfish meatball by adding kelor (*Moringa oleifera* Lam) leaves. International Food Research Journal, 26(1), 263–268.
- More, S. B., Gogate, P. R., & Waghmare, J. S. (2022). Bioactives from pomegranate peel and moringa leaves as natural antioxidants for stability of edible oil blends. Brazilian Journal of Chemical Engineering, 39(2), 527-538.
- Najmah, N., Fitria, R., & Kurniawati, E. (2023). Skrining fitokimia, total flavonoid dan fenolik daun sereh wangi (*Cymbopogon nardus* (L.) Rendle). Jurnal Crystal: Publikasi Penelitian Kimia dan Terapan, 5(1), 62-70 <https://doi.org/10.36526/jc.v5i1.2642>
- Nurfina, N., & Sutji, W. (2021). Pengaruh Penambahan Bahan Fungsional pada Minuman Serbuk Daun Kelor. Prosiding Seminar Nasional Sains dan Teknologi.
- Okoye. (2023). Nutrient and phytochemical compositions of aqueous extracts of black pepper (*Piper guineense*) seed, turmeric (*Curcuma longa*) and ginger (*Zingiber officinale*) rhizome. Journal of Home Economics Research, 107.
- Oluwafemi, A. G., Ajayi, O. B., & Oseni, O. A. (2022). Phytochemical screening, nutritional composition, and antioxidant activities of turmeric (*Curcuma longa*) found in Ado-ekiti, Nigeria. J Appl Life Sci Int, 25(1), 1-8.
- Pan, D., Machado, L., Bica, C. G., Machado, A. K., Steffani, J. A., & Cadoná, F. C. (2022). In vitro evaluation of antioxidant and anticancer activity of lemongrass (*Cymbopogon citratus* (DC) Stapf). Nutrition and cancer, 74(4), 1474-1488. <https://doi.org/10.1080/01635581.2021.1952456>
- Permatananda, P. A. N. K., Aryastuti, A. A. S. A., Cahyawati, P. N., Udiyani, D. P. C., Wijaya, D., Pandit, I. G. S., & Wirajaya, A. A. N. M. (2021, April). Phytochemical and antioxidant capacity test on turmeric extract (*Curcuma longa*) traditionally processed in Bali. In Journal of Physics: Conference Series (Vol. 1869, No. 1, p. 012035). IOP Publishing. <https://doi.org/10.51172/jbmb.v1i2.115>
- Putra, A. I. Y. D., Setiawan, N. B. W., Sanjiwani, M. I. D., Wahyuniari, I. A. I., & Indrayani, A. W. (2021). Nutrigenomic and biomolecular aspect of *Moringa oleifera* leaf powder as supplementation for stunting children. Journal of Tropical Biodiversity and Biotechnology, 6(1), 1-15.

- Putri, A. N., Suryani, I., & Ratnasari, D. (2019). Analisis Uji Sensori untuk Pengembangan Produk Minuman Fungsional Berbasis Bahan Alami. *Jurnal Teknologi Pangan*, 8(2), 102-109.
- Ramanunny, A. K., Wadhwa, S., Gulati, M., Vishwas, S., Khursheed, R., Paudel, K. R., ... & Singh, S. K. (2022). Journey of *Alpinia galanga* from kitchen spice to nutraceutical to folk medicine to nanomedicine. *Journal of ethnopharmacology*, 291, 115144. <https://doi.org/10.1016/j.jep.2022.115144>
- Raczyńska, A., Leszczyńska, T., Skotnicki, P., & Koronowicz, A. (2025). The impact of selected components used in clinical nutrition on inflammation caused malnutrition-a literature review.
- Saepudin, M. S., Pd, M., & Ilyas, M. A. (2024). *Institusi Islam dan pengendalian stunting*. Kbm Kreator Yogyakarta.
- Saini RK, Manoj P, Shetty NP et al. Dietary iron supplements and *Moringa oleifera* leaves influence the liver hepcidin messenger RNA expression and bio-chemical indices of iron status in rats. *Nutr Res* 2014;34:630–8. Crossref. PubMed. <https://doi.org/10.1016/j.nutres.2014.07.003>
- Santoso, A. L., Sanjaya, A., Khamidah, N., & Idawati, I. (2024, January). ENVIRONMENTAL ENTERIC DYSFUNCTION PADA STUNTING. In *Prosiding Seminar Nasional COSMIC Kedokteran* (Vol. 2, pp. 8-12). <https://prosidingcosmic.fk.uwks.ac.id>
- Sharma, S., Katoch, V., Kumar, S., & Chatterjee, S. (2021). Functional relationship of vegetable colors and bioactive compounds: Implications in human health. *The Journal of Nutritional Biochemistry*, 92, 108615. <https://doi.org/10.1016/j.jnutbio.2021.108615>
- Shija AE, Rumisha SF, Oriyo NM et al. Effect of *Moringa oleifera* leaf powder supplementation on reducing anemia in children below two years in Kisarawe District, Tanzania. *Food Sci Nutr* 2019;7:2584–94. Crossref. PubMed. <https://doi.org/10.1002/fsn3.1110>
- Sianipar, E. A. (2021). The potential of Indonesian traditional herbal medicine as immunomodulatory agents: a re-view. *International Journal of Pharmaceutical Sciences and Re-search*, 12(10), 5229. [http://dx.doi.org/10.13040/IJPSR.0975-8232.12\(10\).5229-37](http://dx.doi.org/10.13040/IJPSR.0975-8232.12(10).5229-37)
- Singh, D. N., Bohra, J. S., Dubey, T. P., Shivahre, P. R., Singh, R. K., Singh, T., & Jaiswal, D. K. (2023). Common foods for boosting human immunity: A review. *Food science & nutrition*, 11(11), 6761-6774. <https://doi.org/10.1002/fsn3.3628>
- Su, B., & Chen, X. (2020). Current status and potential of *Moringa oleifera* leaf as an alternative protein source for animal feeds. *Frontiers in veterinary science*, 7, 53.
- Swari, S. J., Sasmita, I. R. A., Kartika, R. C., Alfiansyah, G., & Pratama, M. R. *Produk Inovasi Es Krim Kelor Sebagai Upaya Pencegahan Stunting di Desa*. National Conference for Community Service (NaCosVi), 360–365; 2023 <https://proceedings.polije.ac.id/index.php/ppm/article/view/632>
- Syafitri, N. E., Bintang, M. dan Falah, S. 2014. Kandungan fitokimia, total fenol, dan total flavonoid ekstrak buah harendong (*Melastoma affine* D. Don). *Current Biochemistry*, 1(3), pp.105-115.
- Unuofin, J. O., Masuku, N. P., Paimo, O. K., & Lebelo, S. L. (2021). Ginger from farmyard to town: Nutritional and pharmacological applications. *Frontiers in Pharmacology*, 12, 779352. <https://doi.org/10.3389/fphar.2021.779352>
- Verma, R. K., & Sharma, N. (2022). Phytochemical and pharmacological activities of: a review *Alpinia galanga*. *Asian J Pharm Pharmacol*, 8(3), 74-85.
- Wibowo, A., & Hidayat, T. (2020). Peran Evaluasi Sensoris dalam Optimasi Produk Minuman Sehat. *Prosiding Seminar Nasional Pengembangan Pangan dan Gizi*.
- Wijanarko, A., Perawati, S. dan Andriani, L. 2020. Standardisasi simplisia daun ciplukan. *Jurnal Farmasetis*, 9(1),

- pp.31-40.
<https://doi.org/10.32583/farmasetis.v9i1.736>
- Wijaya, H., Novitasari. dan Jubaidah, S. 2018. Perbandingan metode ekstraksi terhadap rendemen ekstrak daun rambai laut (*Sonneratia caseolaris* L. Engl). *Jurnal Ilmiah Manuntung*, 4(1), pp.79-83.
- Zakaria, N., Fauziah, F., Rinaldi, R., Safrida, Y. D., & Winingsih, E. (2023). Analisa Fitokimia dan Aktivitas Antioksidan Sirup Polih herbal yang Mengandung Daun Kelor, Rimpang Kunyit, dan Rimpang Jahe. *Jurnal Sains dan Kesehatan*, 5(1), 14-21.
- Zhang, J., Xu, X., Cao, Z., Wang, Y., Yang, H., Azarfar, A., & Li, S. (2019). Effect of different tannin sources on nutrient intake, digestibility, performance, nitrogen utilization, and blood parameters in dairy cows. *Animals*, 9(8), 507. <https://doi.org/10.3390/ani9080507>
- Zhang, Y., Hao, R., Chen, J., Li, S., Huang, K., Cao, H., Farag, M. A., Battino, M., Daglia, M., & Capanoglu, E. (2023). Health benefits of saponins and its mechanisms: perspectives from absorption, metabolism, and interaction with gut. *Critical Reviews in Food Science and Nutrition*, 1–22. <https://doi.org/10.1080/10408398.2023.2212063>